

Chiral Symmetry Amplitudes in the S-Wave Isoscalar and Isovector Channels and the σ , $f_0(980)$, $a_0(980)$ SCALAR MESONS

J.A. Oller and E. Oset

*Departamento de Física Teórica and IFIC
Centro Mixto Universidad de Valencia-CSIC
46100 Burjassot (Valencia), Spain*

Abstract. We use a nonperturbative approach which combines coupled channel Lippmann Schwinger equations with meson-meson potentials provided by the lowest order chiral Lagrangian. By means of one parameter, a cut off in the momentum of the loop integrals, which results of the order of 1 GeV, we obtain singularities in the S-wave amplitudes corresponding to the σ , f_0 and a_0 resonances. The $\pi\pi \rightarrow \pi\pi$, $\pi\pi \rightarrow K\bar{K}$ phase shifts and inelasticities in the $T = 0$ scalar channel are well reproduced as well as the $\pi^0\eta$ and $K\bar{K}$ mass distributions in the $T = 1$ channel. Furthermore, the total and partial decay widths of the f_0 and a_0 resonances are properly reproduced. The results seem to indicate that chiral symmetry constraints at low energy and unitarity in coupled channels is the basic information contained in the meson-meson interaction below $\sqrt{s} \simeq 1.2 \text{ GeV}$.

I INTRODUCTION

The understanding of the meson-meson interaction in the scalar sector is still problematic. There is some debate about the spectrum of hadronic states and even more about its nature. Below $\sqrt{s} = 1.2 \text{ GeV}$, which we will study here, the existence of a broad scalar-isoscalar meson around 500 MeV has had permanent ups and downs. However, the $f_0(980)$ (also called S^* , $I^G(J^{PC}) = 0^+(0^{++})$), and the $a_0(980)$ (also called δ , $I^G(J^{PC}) = 1^-(0^{++})$) mesonic states are well established experimentally, although there is still debate around their decay widths and particularly about their nature.

II $L = 0, T = 0, 1$ STRONG AMPLITUDES IN THE COUPLED CHANNEL APPROACH

We assume that the lowest order Hamiltonian provides us with the potential that we iterate in the Lippmann Schwinger equation with two coupled channels, using relativistic meson propagators in the intermediate states. Our channels are labelled 1 for the $K\bar{K}$ and 2 for the $\pi\pi$ states in $T = 0$ and 1 for $K\bar{K}$, 2 for $\pi\eta$ in $T = 1$.

$$\mathcal{L}_2 = \frac{1}{12f^2} < (\partial_\mu \Phi \Phi - \Phi \partial_\mu \Phi)^2 + M\Phi^4 > \quad (1)$$

$$t_{ij} = V_{ij} + \sum_k V_{ik} G_{kk} t_{kj} \quad G_{ii} = i \frac{1}{q^2 - m_{1i}^2 + i\epsilon} \frac{1}{(P-q)^2 - m_{2i}^2 + i\epsilon} \quad (2)$$

where P is the total fourmomentum of the meson-meson systems and q the fourmomentum of one intermediate meson. The former equation is, of course, an integral one. The loop integral in eq. (2) is divergent. In our approach we take a cut off q_{max} ($\Lambda = \sqrt{m_k^2 + q_{max}^2}$) for the maximum value of the modulus of the momentum q . On grounds of chiral symmetry computations we should expect this cut off to be around 1 GeV [1]. By fixing the cut off one generates the counterterms in our scheme.

Let us see that only on shell information is needed. The argument goes as follows: We can write the off shell amplitudes coming from $\mathcal{L}_2$ as

$$V = V_{on} + \beta \sum_i (p_i^2 - m_i^2) \quad V^2 = V_{on}^2 + 2\beta V_{on} \sum_i (p_i^2 - m_i^2) + \beta^2 \sum_{ij} (p_i^2 - m_i^2)(p_j^2 - m_j^2) \quad (3)$$

When performing the q^0 integration in the loop we have two poles. Let us take the contribution from the first pole (the procedure follows analogously for the second pole). From the second term in eq. (3) we get the contribution

$$\frac{2\beta V_{on}}{(2\pi)^3} \int \frac{d^3 q}{2w_1(q)} \frac{(P^0 - w_1)^2 - w_2^2}{(P^0 - w_1)^2 - w_2^2} = \frac{\beta V_{on}}{2\pi^2} \int dw_1 q \quad (4)$$

which for large Λ compared to the masses goes as $V_{on}\Lambda^2$ and has the same structure in the dynamical variables as the tree diagram. The same happens for the rest of off shell terms, that is, they combine with the tree level contribution, giving rise to an amplitude with the same structure as the tree level one but with renormalised parameters f and masses. However, since we are taking physical values for f ($f = f_\pi = 93 \text{ MeV}$) and the masses in the potential, these terms should be omitted. One can proceed like that to higher orders with the same conclusions. Since we are taking V and t on shell they factorize outside the q integral. Thus the term VGT, after the q^0 integration is performed, is given by

$$V_{ij}G_{jj}t_{jk} = V_{ij}(s)t_{jk}(s)G_{jj}(s) \quad (5)$$

$$G_{jj}(s) = \int_0^{q_{max}} \frac{q^2 dq}{(2\pi)^2} \frac{\omega_1 + \omega_2}{\omega_1 \omega_2 [P^{02} - (\omega_1 + \omega_2)^2 + i\epsilon]}$$

where $\omega_i = (\vec{q}^2 + m_i^2)^{1/2}$ and $P^{02} = s$ and the subindex i stands for the two intermediate mesons of the j channel. Thus the coupled channel Lippmann Schwinger equations get reduced to a set of algebraic equations which are solved in a trivial way. Note also that $V_{12} = V_{21}$ and $t_{12} = t_{21}$ by time reversal invariance.

$$\begin{aligned} \Delta_\pi &= 1 - V_{22}G_{22} & \Delta_K &= 1 - V_{11}G_{11} \\ \Delta_c &= \Delta_K \Delta_\pi - V_{12}^2 G_{11} G_{22} & \Delta &= \det A = \Delta_\pi \Delta_c \end{aligned} \quad (6)$$

$$\begin{aligned} t_{11} &= \frac{1}{\Delta_c} (\Delta_\pi V_{11} + V_{12}^2 G_{22}) & t_{21} &= \frac{1}{\Delta_c} (V_{21} G_{11} V_{11} + \Delta_K V_{21}) \\ t_{22} &= \frac{1}{\Delta_c} (\Delta_K V_{22} + V_{12}^2 G_{11}) \end{aligned} \quad (7)$$

The physical amplitudes as a function of s (real variable) are given by the expressions in eq. (7). However, in order to explore the position of the poles of the scattering amplitudes one must take into account the analytical structure of these amplitudes in the different Riemann sheets. These sheets appear because of the cuts related to the opening of thresholds in $G_{jj}(s)$ (see eq. (5)).

III RESULTS

Table I: f_0 Mass and partial widths
($\Lambda = 1.03 \text{ GeV}$)

f_0	our results	experiment
Mass		
Pole position [MeV]	993.5	980 ± 10
Peak of the t_{21} amplitude [MeV]	982	980 ± 10
Γ_{tot} [MeV]	25	40 - 100
$\frac{\Gamma_{\pi\pi}}{\Gamma_{\pi\pi} + \Gamma_{K\bar{K}}}$	0.68	0.67 ± 0.009 0.81 ± 0.004 0.78 ± 0.003 (av.) 0.781 ± 0.024
$\Gamma_{\gamma\gamma}$ (KeV)	0.2	0.63 ± 0.14 $0.42 \pm 0.06 \pm 0.18$ $0.29 \pm 0.07 \pm 0.12$ $0.31 \pm 0.14 \pm 0.09$ (av.) 0.56 ± 0.11

Table 2: a_0 Mass and partial width
($\Lambda = 1.03 \text{ GeV}$)

a_0	our results	experiment
Mass		
Pole position [MeV]	1009.2	983.5 ± 0.9
Peak of the t_{21} amplitude [MeV]	982	983 ± 0.9
Γ_{tot} [MeV]	112	50 - 100
$\frac{\Gamma_{K\bar{K}}}{\Gamma_{\eta\pi}}$	0.60	1.16 ± 0.18 0.7 ± 0.3 0.25 ± 0.008
$\frac{\Gamma_{\gamma\gamma} \cdot \Gamma_{\eta\pi}}{\Gamma_{tot}}$ [KeV]	0.49	$0.28 \pm 0.04 \pm 0.01$ $0.19 \pm 0.07 \pm 0.07$ av. 0.24 ± 0.007

All these data can be obtained from ref. [2], PDG.

Our calculation for the $\gamma\gamma$ width of both resonances is explained in our other contribution to these proceedings, **THEORETICAL STUDY OF THE $\gamma\gamma \rightarrow \text{MESON-MESON REACTION}$.**

σ pole: Mass :469MeV Width :406MeV




FIGURE 1. $I=0$, phase shifts for $\pi\pi \rightarrow \pi\pi$, $K\bar{K} \rightarrow \pi\pi$ and inelasticities. $I=1$, mass distribution of $\pi^-\eta$ in $K^-p \rightarrow \Sigma^+(1385)\pi^-\eta$

IV CONCLUSIONS

We have used a non perturbative approach to deal with the meson meson interaction in the scalar sector at energies below $\sqrt{s} \simeq 1.2 \text{ GeV}$, exploiting chiral symmetry and unitarity in the coupled channels. The Lippmann Schwinger equation with coupled channels, using relativistic meson propagators, and the lowest order chiral Lagrangians, providing the meson-meson potential, are the two ingredients of the theory. One cut off in momentum is also in order to cut the loop integrals and it is fitted to the data [3]. This is the only parameter of the theory. The best fit is found with $q_{max} \simeq 900 \text{ GeV}$, or equivalently $\Lambda = 1.03 \text{ GeV}$. With only these elements we find a good agreement with the experiment for phase shifts, inelasticities and mass distributions until $\sqrt{s} \simeq 1.2 \text{ GeV}$ generating dynamically the $f_0(980)$, $a_0(980)$ and σ resonances.

REFERENCES

1. A. Manohar and Georgi, Nucl. Phys. B234 (1984) 189.
2. R.M. Barnett et al., Phys. Rev. D54 (1996).
3. J.A. Oller and E. Oset, Nucl. Phys. A620 (1997) 438.